
Search for the beta-decay of ⁵⁴Mn


I. Ahmad**, S. Austin§, S. J. Freedman†‡, K. T. Lesko‡, J. L. Mortara*†, A. Wuosmaa**

The decay of ⁵⁴Mn is well known to proceed by electron capture with a half-life of 312 days (Fig. 1). Although it is energetically possible for ⁵⁴Mn to decay to the ground state of ⁵⁴Fe by decay or to the ground state of 54Cr by +, only upper limits have been set for these second forbidden decay modes.1 The current limit on these branches is 10^6 for the - decay and 10^{-8} for the for the + decay. The difference in the branching fractions arises from the greater available phase space for the - decay. These hindered decay modes are the primary means by which a fully ionized ⁵⁴Mn nucleus would decay. Such fully stripped ions are found in cosmic rays and abundances of 54Mn have been reported.2 These relative abundances, combined with measured partial half-lives, provide a cosmic ray chronometer by which one can infer the cosmic-ray confinement time.



Previous efforts¹ have used the characteristic annihilation radiation from positrons as a signature for the + decay. We intend to use annihilation radiation as a trigger with an apparatus designed specifically for such detection. The APEX spectrometer³ was designed to detect the annihilation radiation of positrons created following heavy ion collisions. APEX is a 3 meter long solenoid which will be used to

transport positrons to a charge particle detector surrounded by a segmented annihilation radiation detector. The method has high efficiency and allows us to use an extremely intense radioactive source without suffer from the potentially large 835 keV gamma-ray background.



Footnotes and References

*P-23 Neutron Science and Technology, Los Alamos National Laboratory

†Department of Physics, University of California at Berkeley

‡Nuclear Science Division, Lawrence Berkeley National Laboratory

** Physics Division, Argonne National Laboratory § Michigan State University

1. M. T. F. da Cruz et. al, Physical Review C, 48, 3110 (1993).

2. R. A. Leske, Astrophys. J. 405, 567 (1993).

3. I. Ahmad et. al., Nuclear Instruments and Methods A 370, 539 (1996).

4. M. DuVernois, private communication.